

SARAFEN-P

Each tablet contains:

Composition:

Serratiopeptidase	15 mg
Paracetamol	500 mg
Aceclofenac	100 mg

Indications:

Rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, cervical spondylitis, intervertabral disc syndrome, sciatica, non-articular rheumatic conditions. Post-operative & traumatic inflammations. Painful inflammatory conditions in gynaecology & dentistry. Acute gout.

Description:

<u>Serratiopeptidase</u> is a proteolytic enzyme derived from serratia Spp. <u>Serratiopeptidase</u> binds to alpha-2 macroglobulin in the blood in the ratio of 1:1 which helps to mask its antigenicity but retain its enzymatic activity by hydrolysing bradykinin, histamine, and serotonin. Serratiopeptidase blocks plasmin inhibition thus helping the fibrinolytic activity of plasmin helps clearance of exudate, reduces swelling and improves micro circulation.

CONTRAINDICATIONS

Blood coagulation abnormalities disturbance or under treatment with anticoagulants.

SIDE-EFFECTS: G.I. disturbance like anorexia, gastric discomfort and nausea. Discontinue if hypersensitive symptoms such as eruption appear. **Dosage:** As per the physician's advice.

Presentations: 10 tablets

 MRP
 Retailer
 Stockiest

 42.80
 34.24
 30.81